



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 39/395, A61P 35/00 // (A61K 39/395, 31:335)	A1	(11) International Publication Number: WO 00/69460 (43) International Publication Date: 23 November 2000 (23.11.00)
(21) International Application Number: PCT/US00/12552 (22) International Filing Date: 9 May 2000 (09.05.00) (30) Priority Data: 60/134,085 14 May 1999 (14.05.99) US (71) Applicant: GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US). (72) Inventor: COHEN, Robert, L.; 660 Parrott Drive, San Mateo, CA 94402 (US). (74) Agents: LEE, Wendy, M. et al.; Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080-4990 (US).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: TREATMENT WITH ANTI-ErbB2 ANTIBODIES (57) Abstract A method treating a human patient to or diagnosed with a tumor in which ErbB2 protein is expressed comprising the following steps, performed sequentially: (a) treating the patient with a therapeutically effective amount of an anti-ErbB2 antibody; (b) surgically removing the tumor, and then (c) treating the patient with a therapeutically effective amount of an anti-ErbB2 antibody or of a chemotherapeutic agent.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TREATMENT WITH ANTI-ErbB2 ANTIBODIES

Field of the Invention

5 The present invention concerns the treatment of cancer with anti-ErbB2 antibodies.

Background of the Invention

Proto-oncogenes that encode growth factors and growth factor receptors have been identified to play important roles in the pathogenesis of various human malignancies, including breast cancer. It has been found that the human *erbB2* gene (also known as *HER2*, or *c-erbB-2*), which encodes a 185-kd transmembrane glycoprotein receptor (p185^{HER2}) related to the epidermal growth factor receptor (EGFR), is overexpressed in about 25% to 30% of human breast cancer (Slamon *et al.*, *Science* 235:177-182 [1987]; Slamon *et al.*, *Science* 244:707-712 [1989]).

Several lines of evidence support a direct role for ErbB2 in the pathogenesis and clinical aggressiveness of ErbB2-overexpressing tumors. The introduction of ErbB2 into non-neoplastic cells has been shown to cause their malignant transformation (Hudziak *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7159-7163 [1987]; DiFiore *et al.*, *Science* 237:78-182 [1987]). Transgenic mice that express HER2 were found to develop mammary tumors (Guy *et al.*, *Proc. Natl. Acad. Sci. USA* 89:10578-10582 [1992]).

Antibodies directed against human ErbB2 protein and the protein encoded by the rat equivalent of the *erbB2* gene (*neu*) have been described. Drebin *et al.*, *Cell* 41:695-706 (1985) refer to an IgG2a monoclonal antibody which is directed against the rat *neu* gene product. This antibody called 7.16.4 causes down-modulation of cell surface p185 expression on B104-1-1 cells (NIH-3T3 cells transfected with the *neu* protooncogene) and inhibits colony formation of these cells. In Drebin *et al.* *PNAS (USA)* 83:9129-9133 (1986), the 7.16.4 antibody was shown to inhibit the tumorigenic growth of *neu*-transformed NIH-3T3 cells as well as rat neuroblastoma cells (from which the *neu* oncogene was initially isolated) implanted into nude mice. Drebin *et al.* in *Oncogene* 2:387-394 (1988) discuss the production of a panel of antibodies against the rat *neu* gene product. All of the antibodies were found to exert a cytostatic effect on the growth of *neu*-transformed cells suspended in soft agar. Antibodies of the IgM, IgG2a and IgG2b isotypes were able to mediate significant *in vitro* lysis of *neu*-transformed cells in the presence of complement, whereas none of the antibodies were able to mediate high levels of antibody dependent cellular cytotoxicity (ADCC) of the *neu*-transformed cells. Drebin *et al.* *Oncogene* 2:273-277 (1988) report that mixtures of antibodies reactive with two distinct regions on the p185 molecule result in synergistic anti-tumor effects on *neu*-transformed NIH-3T3 cells implanted into nude mice. Biological effects of anti-*neu* antibodies are reviewed in Myers *et al.*, *Meth. Enzym.* 198:277-290 (1991). See also WO 94/22478 published October 13, 1994.

Hudziak *et al.*, *Mol. Cell. Biol.* 9(3):1165-1172 (1989) describe the generation of a panel of anti-ErbB2 antibodies which were characterized using the human breast tumor cell line SKBR3. Relative cell proliferation of the SKBR3 cells following exposure to the antibodies was determined by crystal violet staining of the monolayers after 72 hours. Using this assay, maximum inhibition was obtained with the antibody called 4D5 which inhibited cellular proliferation by 56%. Other antibodies in the panel, including 7C2 and 7F3, reduced cellular proliferation to a lesser extent in this assay. Hudziak *et al.* conclude that the effect of the 4D5 antibody on SKBR3 cells was cytostatic rather than cytotoxic, since SKBR3 cells resumed growth at a nearly normal rate following removal of

the antibody from the medium. The antibody 4D5 was further found to sensitize p185^{erbB2}-overexpressing breast tumor cell lines to the cytotoxic effects of TNF- α . See also WO89/06692 published July 27, 1989. The anti-ErbB2 antibodies discussed in Hudziak *et al.* are further characterized in Fendly *et al. Cancer Research* 50:1550-1558 (1990); Kotts *et al. In Vitro* 26(3):59A (1990); Sarup *et al. Growth Regulation* 1:72-82 (1991); Shepard *et al. J. Clin. Immunol.* 11(3):117-127 (1991); Kumar *et al. Mol. Cell. Biol.* 11(2):979-986 (1991); Lewis *et al. Cancer Immunol. Immunother.* 37:255-263 (1993); Pietras *et al. Oncogene* 9:1829-1838 (1994); Vitetta *et al. Cancer Research* 54:5301-5309 (1994); Sliwkowski *et al. J. Biol. Chem.* 269(20):14661-14665 (1994); Scott *et al. J. Biol. Chem.* 266:14300-5 (1991); and D'souza *et al. Proc. Natl. Acad. Sci.* 91:7202-7206 (1994).

Tagliabue *et al. Int. J. Cancer* 47:933-937 (1991) describe two antibodies which were selected for their reactivity on the lung adenocarcinoma cell line (Calu-3) which overexpresses ErbB2. One of the antibodies, called MGR3, was found to internalize, induce phosphorylation of ErbB2, and inhibit tumor cell growth *in vitro*.

McKenzie *et al. Oncogene* 4:543-548 (1989) generated a panel of anti-ErbB2 antibodies with varying epitope specificities, including the antibody designated TA1. This TA1 antibody was found to induce accelerated endocytosis of ErbB2 (see Maier *et al. Cancer Res.* 51:5361-5369 (1991)). Bacus *et al. Molecular Carcinogenesis* 3:350-362 (1990) reported that the TA1 antibody induced maturation of the breast cancer cell lines AU-565 (which overexpresses the *erbB2* gene) and MCF-7 (which does not). Inhibition of growth and acquisition of a mature phenotype in these cells was found to be associated with reduced levels of ErbB2 receptor at the cell surface and transient increased levels in the cytoplasm.

Stancovski *et al. PNAS (USA)* 88:8691-8695 (1991) generated a panel of anti-ErbB2 antibodies, injected them i.p. into nude mice and evaluated their effect on tumor growth of murine fibroblasts transformed by overexpression of the *erbB2* gene. Various levels of tumor inhibition were detected for four of the antibodies, but one of the antibodies (N28) consistently stimulated tumor growth. Monoclonal antibody N28 induced significant phosphorylation of the ErbB2 receptor, whereas the other four antibodies generally displayed low or no phosphorylation-inducing activity. The effect of the anti-ErbB2 antibodies on proliferation of SKBR3 cells was also assessed. In this SKBR3 cell proliferation assay, two of the antibodies (N12 and N29) caused a reduction in cell proliferation relative to control. The ability of the various antibodies to induce cell lysis *in vitro* via complement-dependent cytotoxicity (CDC) and antibody dependent cellular cytotoxicity (ADCC) was assessed, with the authors of this paper concluding that the inhibitory function of the antibodies was not attributed significantly to CDC or ADCC.

Bacus *et al. Cancer Research* 52:2580-2589 (1992) further characterized the antibodies described in Bacus *et al.* (1990) and Stancovski *et al.* of the preceding paragraphs. Extending the i.p. studies of Stancovski *et al.*, the effect of the antibodies after i.v. injection into nude mice harboring mouse fibroblasts overexpressing human ErbB2 was assessed. As observed in their earlier work, N28 accelerated tumor growth whereas N12 and N29 significantly inhibited growth of the ErbB2-expressing cells. Partial tumor inhibition was also observed with the N24 antibody. Bacus *et al.* also tested the ability of the antibodies to promote a mature phenotype in the human breast cancer cell lines AU-565 and MDA-MB453 (which overexpress ErbB2) as well as MCF-7 (containing low levels of the receptor). Bacus *et al.* saw a correlation between tumor inhibition *in vivo* and cellular differentiation;

the tumor-stimulatory antibody N28 had no effect on differentiation, and the tumor inhibitory action of the N12, N29 and N24 antibodies correlated with the extent of differentiation they induced.

Xu *et al.* *Int. J. Cancer* 53:401-408 (1993) evaluated a panel of anti-ErbB2 antibodies for their epitope binding specificities, as well as their ability to inhibit anchorage-independent and anchorage-dependent growth of SKBR3 cells (by individual antibodies and in combinations), modulate cell-surface ErbB2, and inhibit ligand stimulated anchorage-independent growth. See also WO94/00136 published Jan 6, 1994 and Kasprzyk *et al.* *Cancer Research* 52:2771-2776 (1992) concerning anti-ErbB2 antibody combinations. Other anti-ErbB2 antibodies are discussed in Hancock *et al.* *Cancer Res.* 51:4575-4580 (1991); Shawver *et al.* *Cancer Res.* 54:1367-1373 (1994); Arteaga *et al.* *Cancer Res.* 54:3758-3765 (1994); and Harwerth *et al.* *J. Biol. Chem.* 267:15160-15167 (1992).

A recombinant humanized anti-ErbB2 monoclonal antibody (a humanized version of the murine anti-ErbB2 antibody 4D5, referred to as rhuMab HER2 or HERCEPTIN[®]) has been clinically active in patients with ErbB2-overexpressing metastatic breast cancers that had received extensive prior anticancer therapy. (Baselga *et al.*, *J. Clin. Oncol.* 14:737-744 [1996]).

ErbB2 overexpression is commonly regarded as a predictor of a poor prognosis, especially in patients with primary disease that involves axillary lymph nodes (Slamon *et al.*, [1987] and [1989], *supra*; Ravdin and Chamness, *Gene* 159:19-27 [1995]; and Hynes and Stern, *Biochim Biophys Acta* 1198:165-184 [1994]), and has been linked to sensitivity and/or resistance to hormone therapy and chemotherapeutic regimens, including CMF (cyclophosphamide, methotrexate, and fluoruracil) and anthracyclines (Baselga *et al.*, *Oncology* 11(3 Suppl 1):43-48 [1997]). However, despite the association of ErbB2 overexpression with poor prognosis, the odds of HER2-positive patients responding clinically to treatment with taxanes were greater than three times those of HER2-negative patients (*Ibid*). rhuMab HER2 was shown to enhance the activity of paclitaxel (TAXOL[®]) and doxorubicin against breast cancer xenografts in nude mice injected with BT-474 human breast adenocarcinoma cells, which express high levels of HER2 (Baselga *et al.*, *Breast Cancer, Proceedings of ASCO*, Vol. 13, Abstract 53 [1994]).

Summary of the Invention

In a first aspect, the present invention provides a method of treating a human patient susceptible to or diagnosed with a tumor in which ErbB2 protein is expressed comprising the following steps, performed sequentially:

- (a) treating the patient with a therapeutically effective amount of an anti-ErbB2 antibody and, optionally, further comprising treating the patient with a therapeutically effective amount of a chemotherapeutic agent (e.g. a taxoid, such as paclitaxel or doxetaxel);
- (b) surgically removing the tumor; and then
- (c) treating the patient with a therapeutically effective amount of an anti-ErbB2 antibody and/or of a chemotherapeutic agent (e.g. a taxoid, such as paclitaxel or doxetaxel).

Preferably, the tumor overexpresses ErbB2 protein and is selected from the group consisting of a breast tumor, squamous cell tumor, small-cell lung tumor, non-small cell lung tumor, gastrointestinal tumor, pancreatic tumor, glioblastoma, cervical tumor, ovarian tumor, liver tumor, bladder tumor, hepatoma, colon tumor, colorectal

tumor, endometrial tumor, salivary gland tumor, kidney tumor, prostate tumor, vulval tumor, thyroid tumor, hepatic tumor, head tumor and neck tumor.

The invention further provides an article of manufacture comprising a container, a composition within the container comprising an anti-ErbB2 antibody and a package insert instructing the user of the composition to treat a patient essentially according to the above method.

Brief Description of the Drawings

Fig. 1 shows epitope-mapping of the extracellular domain of ErbB2 as determined by truncation mutant analysis and site-directed mutagenesis (Nakamura *et al. J. of Virology* 67(10):6179-6191 [Oct 1993]; Renz *et al. J. Cell Biol.* 125(6):1395-1406 [Jun 1994]). The anti-proliferative MABs 4D5 and 3H4 bind adjacent to the transmembrane domain. The various ErbB2-ECD truncations or point mutations were prepared from cDNA using polymerase chain reaction technology. The ErbB2 mutants were expressed as gD fusion proteins in a mammalian expression plasmid. This expression plasmid uses the cytomegalovirus promoter/enhancer with SV40 termination and polyadenylation signals located downstream of the inserted cDNA. Plasmid DNA was transfected into 293S cells. One day following transfection, the cells were metabolically labeled overnight in methionine and cysteine-free, low glucose DMEM containing 1% dialyzed fetal bovine serum and 25 μ Ci each of 35 S methionine and 35 S cysteine. Supernatants were harvested either the ErbB2 MABs or control antibodies were added to the supernatant and incubated 2-4 hours at 4°C. The complexes were precipitated, applied to a 10-20% Tricine SDS gradient gel and electrophoresed at 100 V. The gel was electroblotted onto a membrane and analyzed by autoradiography. SEQ ID NO:3 and 4 depict the 3H4 and 4D5 epitopes, respectively.

Fig. 2 depicts with underlining the amino acid sequence of Domain 1 of ErbB2 (SEQ ID NO:1). Bold amino acids indicate the location of the epitope recognized by MABs 7C2 and 7F3 as determined by deletion mapping, i.e. the "7C2/7F3 epitope" (SEQ ID NO:2).

Detailed Description of the Preferred Embodiments

L. Definitions

The terms "HER2", "ErbB2" and "c-Erb-B2" are used interchangeably. Unless indicated otherwise, the terms "ErbB2", "c-Erb-B2" and "HER2" refer to the human protein, and "*Her2*", "*erbB2*" and "*c-erb-B2*" refer to human gene. The human *erbB2* gene and ErbB2 protein are, for example, described in Semba *et al., PNAS (USA)* 82:6497-6501 (1985) and Yamamoto *et al. Nature* 319:230-234 (1986) (Genebank accession number X03363). ErbB2 comprises four domains (Domains 1-4).

The "epitope 4D5" is the region in the extracellular domain of ErbB2 to which the antibody 4D5 (ATCC CRL 10463) binds. This epitope is close to the transmembrane region of ErbB2. To screen for antibodies which bind to the 4D5 epitope, a routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. Alternatively, epitope mapping can be performed (see Fig. 1) to assess whether the antibody binds to the 4D5 epitope of ErbB2 (i.e. any one or more residues in the region from about residue 529, e.g. about residue 561 to about residue 625, inclusive; SEQ ID NO:4).

The "epitope 3H4" is the region in the extracellular domain of ErbB2 to which the antibody 3H4 binds. This epitope is shown in Fig. 1, and includes residues from about 541 to about 599, inclusive, in the amino acid sequence of ErbB2 extracellular domain (SEQ ID NO:3).

The "epitope 7C2/7F3" is the region at the N terminus of the extracellular domain of ErbB2 to which the 7C2 and/or 7F3 antibodies (each deposited with the ATCC, see below) bind. To screen for antibodies which bind to the 7C2/7F3 epitope, a routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. Alternatively, epitope mapping can be performed to establish whether the antibody binds to the 7C2/7F3 epitope on ErbB2 (*i.e.* any one or more of residues in the region from about residue 22 to about residue 53 of ErbB2 [SEQ ID NO:2]).

The term "induces cell death" or "capable of inducing cell death" refers to the ability of the antibody to make a viable cell become nonviable. The "cell" here is one which expresses the ErbB2 receptor, especially where the cell overexpresses the ErbB2 receptor. A cell which "overexpresses" ErbB2 has significantly higher than normal ErbB2 levels compared to a noncancerous cell of the same tissue type. Preferably, the cell is a cancer cell, *e.g.* a breast, ovarian, stomach, endometrial, salivary gland, lung, kidney, colon, thyroid, pancreatic or bladder cell. *In vitro*, the cell may be a SKBR3, BT474, Calu 3, MDA-MB-453, MDA-MB-361 or SKOV3 cell. Cell death *in vitro* may be determined in the absence of complement and immune effector cells to distinguish cell death induced by antibody dependent cellular cytotoxicity (ADCC) or complement dependent cytotoxicity (CDC). Thus, the assay for cell death may be performed using heat inactivated serum (*i.e.* in the absence of complement) and in the absence of immune effector cells. To determine whether the antibody is able to induce cell death, loss of membrane integrity as evaluated by uptake of propidium iodide (PI), trypan blue (see Moore *et al.* *Cytotechnology* 17:1-11 [1995]) or 7AAD can be assessed relative to untreated cells. Preferred cell death-inducing antibodies are those which induce PI uptake in the "PI uptake assay in BT474 cells".

The phrase "induces apoptosis" or "capable of inducing apoptosis" refers to the ability of the antibody to induce programmed cell death as determined by binding of annexin V, fragmentation of DNA, cell shrinkage, dilation of endoplasmic reticulum, cell fragmentation, and/or formation of membrane vesicles (called apoptotic bodies). The cell is one which overexpresses the ErbB2 receptor. Preferably the "cell" is a tumor cell, *e.g.* a breast, ovarian, stomach, endometrial, salivary gland, lung, kidney, colon, thyroid, pancreatic or bladder cell. *In vitro*, the cell may be a SKBR3, BT474, Calu 3 cell, MDA-MB-453, MDA-MB-361 or SKOV3 cell. Various methods are available for evaluating the cellular events associated with apoptosis. For example, phosphatidyl serine (PS) translocation can be measured by annexin binding; DNA fragmentation can be evaluated through DNA laddering as disclosed in the example herein; and nuclear/chromatin condensation along with DNA fragmentation can be evaluated by any increase in hypodiploid cells. Preferably, the antibody which induces apoptosis is one which results in about 2 to 50 fold, preferably about 5 to 50 fold, and most preferably about 10 to 50 fold, induction of annexin binding relative to untreated cell in an "annexin binding assay using BT474 cells" (see below).

Sometimes the pro-apoptotic antibody will be one which blocks HRG binding/activation of the ErbB2/ErbB3 complex (*e.g.* 7F3 antibody). In other situations, the antibody is one which does not significantly block activation of the ErbB2/ErbB3 receptor complex by HRG (*e.g.* 7C2). Further, the antibody may be one like

7C2 which, while inducing apoptosis, does not induce a large reduction in the percent of cells in S phase (e.g. one which only induces about 0-10% reduction in the percent of these cells relative to control).

The antibody of interest may be one like 7C2 which binds specifically to human ErbB2 and does not significantly cross-react with other proteins such as those encoded by the *erbB1*, *erbB3* and/or *erbB4* genes. Sometimes, the antibody may not significantly cross-react with the rat *neu* protein, e.g., as described in Schecter *et al.* *Nature* 312:513 (1984) and Drebin *et al.*, *Nature* 312:545-548 (1984). In such embodiments, the extent of binding of the antibody to these proteins (e.g., cell surface binding to endogenous receptor) will be less than about 10% as determined by fluorescence activated cell sorting (FACS) analysis or radioimmunoprecipitation (RIA).

"Heregulin" (HRG) when used herein refers to a polypeptide which activates the ErbB2-ErbB3 and ErbB2-ErbB4 protein complexes (i.e. induces phosphorylation of tyrosine residues in the complex upon binding thereto). Various heregulin polypeptides encompassed by this term are disclosed in Holmes *et al.*, *Science*, 256:1205-1210 (1992); WO 92/20798; Wen *et al.*, *Mol. Cell. Biol.*, 14(3):1909-1919 (1994); and Marchionni *et al.*, *Nature*, 362:312-318 (1993), for example. The term includes biologically active fragments and/or variants of a naturally occurring HRG polypeptide, such as an EGF-like domain fragment thereof (e.g. HRG β 1₁₇₇₋₂₄₄).

The "ErbB2-ErbB3 protein complex" and "ErbB2-ErbB4 protein complex" are noncovalently associated oligomers of the ErbB2 receptor and the ErbB3 receptor or ErbB4 receptor, respectively. The complexes form when a cell expressing both of these receptors is exposed to HRG and can be isolated by immunoprecipitation and analyzed by SDS-PAGE as described in Sliwkowski *et al.*, *J. Biol. Chem.*, 269(20):14661-14665 (1994).

"Antibodies" (Abs) and "immunoglobulins" (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to a specific antigen, immunoglobulins include both antibodies and other antibody-like molecules which lack antigen specificity. Polypeptides of the latter kind are, for example, produced at low levels by the lymph system and at increased levels by myelomas.

"Native antibodies" and "native immunoglobulins" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light-chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light- and heavy-chain variable domains.

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called complementarity determining regions (CDRs) or hypervariable regions both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework region (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and

in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, *NIH Publ. No. 91-3242*, Vol. I, pages 647-669 [1991]). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an $F(ab')_2$ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V_H - V_L dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. $F(ab')_2$ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), *e.g.*, IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy-chain constant domains that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

The term "antibody" is used in the broadest sense and specifically covers intact monoclonal antibodies, polyclonal antibodies, multispecific antibodies (*e.g.* bispecific antibodies) formed from at least two intact antibodies, and antibody fragments so long as they exhibit the desired biological activity.

"Antibody fragments" comprise a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', $F(ab')_2$, and Fv fragments; diabodies; linear antibodies (Zapata *et al. Protein Eng.* 8(10):1057-1062 [1995]); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (see, *e.g.*, U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991), for example.

The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; Morrison *et al.*, *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1984)).

"Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementarity-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and maximize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, *Nature*, 321:522-525 (1986); Reichmann *et al.*, *Nature*, 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992). The humanized antibody includes a PRIMATIZED™ antibody wherein the antigen-binding

region of the antibody is derived from an antibody produced by immunizing macaque monkeys with the antigen of interest.

"Single-chain Fv" or "sFv" antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv see Plückthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) in the same polypeptide chain ($V_H - V_L$). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (*e.g.*, IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule.

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disorder as well as those in which the disorder is to be prevented.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, *etc.* Preferably, the mammal is human.

A "disorder" is any condition that would benefit from treatment with the anti-ErbB2 antibody. This includes chronic and acute disorders or diseases including those pathological conditions which predispose the mammal to the disorder in question. Non-limiting examples of disorders to be treated herein include benign and malignant tumors; leukemias and lymphoid malignancies; neuronal, glial, astrocytal, hypothalamic and other glandular, macrophagal, epithelial, stromal and blastocoelic disorders; and inflammatory, angiogenic and immunologic disorders.

The term "therapeutically effective amount" refers to an amount of a drug effective to treat a disease or disorder in a mammal. In the case of cancer, the therapeutically effective amount of the drug may reduce the number of cancer cells; reduce the tumor size; inhibit (*i.e.*, slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit (*i.e.*, slow to some extent and preferably stop) tumor metastasis; inhibit, to some extent, tumor growth; and/or relieve to some extent one or more of the symptoms associated with the disorder. To the extent the drug may prevent growth and/or kill existing cancer cells, it may be cytostatic and/or cytotoxic. For cancer therapy, efficacy can, for example, be measured by assessing the time to tumor progression (TTP), determining the response rate (RR) and/or evaluating overall survival.

The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include, but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia or lymphoid malignancies. More particular examples of such cancers include squamous cell cancer, lung cancer including small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung and squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer including gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, liver cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma as well as head and neck cancer.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (*e.g.* I^{131} , I^{123} , I^{125} , Y^{90} , At^{211} , Cu^{67} , Bi^{212} , Pd^{109} , Re^{188} and Re^{186}), chemotherapeutic agents, and toxins such as enzymatically active toxins of bacterial, fungal, plant or animal origin, or fragments thereof.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXANTM); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramidate, triethylenethiophosphoramidate and trimethylolomelamine; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, 5-FU; androgens such as calusterone, dromostanolone propionate, epitostanol, mepitostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide

glycoside; aminolevulinic acid; amsacrine; bestabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfomithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK®; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2, 2',2''-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, *e.g.* paclitaxel (TAXOL®, Bristol-Myers Squibb Oncology, Princeton, NJ) and doxetaxel (Taxotere, Rhône-Poulenc Rorer, Antony, France); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; carminomycin; aminopterin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition are hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone; and anti-androgens such as flutamide and nilutamide; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

A "growth inhibitory agent" when used herein refers to a compound or composition which inhibits growth of a cell, especially an ErbB2-overexpressing cancer cell either *in vitro* or *in vivo*. Thus, the growth inhibitory agent is one which significantly reduces the percentage of ErbB2 overexpressing cells in S phase. Examples of growth inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), taxol, and topo II inhibitors such as doxorubicin, epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in *The Molecular Basis of Cancer*, Mendelsohn and Israel, eds., Chapter 1, entitled "Cell cycle regulation, oncogens, and antineoplastic drugs" by Murakami *et al.* (WB Saunders: Philadelphia, 1995), especially p. 13. The 4D5 antibody (and functional equivalents thereof) can also be employed for this purpose.

"Doxorubicin" is an anthracycline antibiotic. The full chemical name of doxorubicin is (8S-cis)-10-[(3-amino-2,3,6-trideoxy- α -L-lyxo-hexapyranosyl)oxy]-7,8,9,10-tetrahydro-6,8,11-trihydroxy-8-(hydroxyacetyl)-1-methoxy-5,12-naphthacenedione.

The term "cytokine" is a generic term for proteins released by one cell population which act on another cell as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor- α and - β ; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve

growth factors such as NGF- β ; platelet-growth factor; transforming growth factors (TGFs) such as TGF- α and TGF- β ; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon- α , - β , and - γ ; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-1 α , IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-11, IL-12; a tumor necrosis factor such as TNF- α or TNF- β ; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

The term "prodrug" as used in this application refers to a precursor or derivative form of a pharmaceutically active substance that is less cytotoxic to tumor cells compared to the parent drug and is capable of being enzymatically activated or converted into the more active parent form. See, e.g., Wilman, "Prodrugs in Cancer Chemotherapy" *Biochemical Society Transactions*, 14, pp. 375-382, 615th Meeting Belfast (1986) and Stella *et al.*, "Prodrugs: A Chemical Approach to Targeted Drug Delivery," *Directed Drug Delivery*, Borchardt *et al.*, (ed.), pp. 247-267, Humana Press (1985). The prodrugs of this invention include, but are not limited to, phosphate-containing prodrugs, thiophosphate-containing prodrugs, sulfate-containing prodrugs, peptide-containing prodrugs, D-amino acid-modified prodrugs, glycosylated prodrugs, β -lactam-containing prodrugs, optionally substituted phenoxyacetamide-containing prodrugs or optionally substituted phenylacetamide-containing prodrugs, 5-fluorocytosine and other 5-fluorouridine prodrugs which can be converted into the more active cytotoxic free drug. Examples of cytotoxic drugs that can be derivatized into a prodrug form for use in this invention include, but are not limited to, those chemotherapeutic agents described above.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as the anti-ErbB2 antibodies disclosed herein and, optionally, a chemotherapeutic agent) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

The term "package insert" is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, contraindications and/or warnings concerning the use of such therapeutic products.

A "cardioprotectant" is a compound or composition which prevents or reduces myocardial dysfunction (*i.e.* cardiomyopathy and/or congestive heart failure) associated with administration of a drug, such as an anthracycline antibiotic and/or an anti-ErbB2 antibody, to a patient. The cardioprotectant may, for example, block or reduce a free-radical-mediated cardiotoxic effect and/or prevent or reduce oxidative-stress injury. Examples of cardioprotectants encompassed by the present definition include the iron-chelating agent dexrazoxane (ICRF-187) (Seifert *et al.* *The Annals of Pharmacotherapy* 28:1063-1072 [1994]); a lipid-lowering agent and/or anti-oxidant such as probucol (Singal *et al.* *J. Mol. Cell Cardiol.* 27:1055-1063 [1995]); amifostine (aminothiols 2-[(3-aminopropyl)amino]ethanethiol-dihydrogen phosphate ester, also called WR-2721, and the dephosphorylated cellular uptake form thereof called WR-1065) and S-3-(3-methylaminopropylamino)propylphosphorothioic acid (WR-151327), see Green *et al.* *Cancer Research* 54:738-741 (1994); digoxin (Bristow, M.R. In: Bristow MR, ed. *Drug-Induced Heart Disease*. New York: Elsevier 191-215 [1980]); beta-blockers such as metoprolol (Hjalmarson *et al.* *Drugs* 47:Suppl 4:31-9 [1994]; and Shaddy *et al.* *Am. Heart J.* 129:197-9 [1995]); vitamin E; ascorbic acid

(vitamin C); free radical scavengers such as oleanolic acid, ursolic acid and N-acetylcysteine (NAC); spin trapping compounds such as alpha-phenyl-tert-butyl nitron (PBN); (Paracchini *et al.*, *Anticancer Res.* 13:1607-1612 [1993]); selenoorganic compounds such as P251 (Elbesen); and the like.

II. Production of anti-ErbB2 antibodies

5 A description follows as to exemplary techniques for the production of the antibodies used in accordance with the present invention. The ErbB2 antigen to be used for production of antibodies may be, *e.g.*, a soluble form of the extracellular domain of ErbB2 or a portion thereof, containing the desired epitope. Alternatively, cells expressing ErbB2 at their cell surface [*e.g.* NIH-3T3 cells transformed to overexpress ErbB2; or a carcinoma cell line such as SKBR3 cells, see Stancovski *et al.* *PNAS (USA)* 88:8691-8695 (1991)] can be used
10 to generate antibodies. Other forms of ErbB2 useful for generating antibodies will be apparent to those skilled in the art.

(i) Polyclonal antibodies

Polyclonal antibodies are preferably raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen and an adjuvant. It may be useful to conjugate the relevant antigen to a protein
15 that is immunogenic in the species to be immunized, *e.g.*, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N}=\text{C}=\text{NR}$, where R and R^1 are different alkyl groups.

20 Animals are immunized against the antigen, immunogenic conjugates, or derivatives by combining, *e.g.*, 100 μg or 5 μg of the protein or conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of peptide or conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites. Seven to 14 days later the animals are bled and the serum is assayed for antibody titer. Animals
25 are boosted until the titer plateaus. Preferably, the animal is boosted with the conjugate of the same antigen, but conjugated to a different protein and/or through a different cross-linking reagent. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are suitably used to enhance the immune response.

(ii) Monoclonal antibodies

30 Monoclonal antibodies are obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Thus, the modifier "monoclonal" indicates the character of the antibody as not being a mixture of discrete antibodies.

For example, the monoclonal antibodies may be made using the hybridoma method first described by
35 Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567).

In the hybridoma method, a mouse or other appropriate host animal, such as a hamster, is immunized as hereinabove described to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*.

Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 [Academic Press, 1986]).

The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

Preferred myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Maryland USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)].

Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA).

The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson *et al.*, *Anal. Biochem.*, 107:220 (1980).

After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 [Academic Press, 1986]). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal.

The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra *et al.*, *Curr. Opinion in Immunol.*, 5:256-262 (1993) and Plückthun, *Immunol. Revs.*, 130:151-188 (1992).

In a further embodiment, antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in McCafferty *et al.*, *Nature*, 348:552-554 (1990). Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991) describe the isolation of murine and human antibodies, respectively, using phage libraries. Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Marks *et al.*, *Bio/Technology*, 10:779-783 [1992]), as well as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage libraries (Waterhouse *et al.*, *Nuc. Acids. Res.*, 21:2265-2266 [1993]). Thus, these techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

The DNA also may be modified, for example, by substituting the coding sequence for human heavy- and light-chain constant domains in place of the homologous murine sequences [U.S. Patent No. 4,816,567; Morrison, *et al.*, *Proc. Natl Acad. Sci. USA*, 81:6851 (1984)], or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide.

Typically such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody, or they are substituted for the variable domains of one antigen-combining site of an antibody to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for an antigen and another antigen-combining site having specificity for a different antigen.

(iii) *Humanized and human antibodies*

Methods for humanizing non-human antibodies are well known in the art. Preferably, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones *et al.*, *Nature*, 321:522-525 (1986); Riechmann *et al.*, *Nature*, 332:323-327 (1988); Verhoeven *et al.*, *Science*, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567) wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework for the humanized antibody [Sims *et al.*, *J. Immunol.*, 151:2296 (1993); Chothia *et al.*, *J. Mol. Biol.*, 196:901 (1987)]. Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies [Carter *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta *et al.*, *J. Immunol.*, 151:2623 (1993)].

It is further important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies

are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *i.e.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding.

Alternatively, it is now possible to produce transgenic animals (*e.g.*, mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, *e.g.*, Jakobovits *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:2551 (1993); Jakobovits *et al.*, *Nature*, 362:255-258 (1993); Bruggermann *et al.*, *Year in Immuno.*, 7:33 (1993). Human antibodies can also be derived from phage-display libraries (Hoogenboom *et al.*, *J. Mol. Biol.*, 227:381 (1991); Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991)).

(iv) Antibody fragments

Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies [see, *e.g.*, Morimoto *et al.*, *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992) and Brennan *et al.*, *Science*, 229:81 (1985)]. However, these fragments can now be produced directly by recombinant host cells. For example, the antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form $F(ab')_2$ fragments [Carter *et al.*, *Bio/Technology* 10:163-167 (1992)]. According to another approach, $F(ab')_2$ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv). See WO 93/16185.

(v) Bispecific antibodies

Bispecific antibodies are antibodies that have binding specificities for at least two different epitopes. Exemplary bispecific antibodies may bind to two different epitopes of the ErbB2 protein. For example, one arm may bind an epitope in Domain I of ErbB2 such as the 7C2/7F3 epitope, the other may bind a different ErbB2 epitope, *e.g.* the 4D5 epitope. Other such antibodies may combine an ErbB2 binding site with binding site(s) for EGFR, ErbB3 and/or ErbB4. Alternatively, an anti-ErbB2 arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (*e.g.* CD2 or CD3), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the ErbB2-expressing cell. Bispecific antibodies may also be used to localize cytotoxic agents to cells which

express ErbB2. These antibodies possess an ErbB2-binding arm and an arm which binds the cytotoxic agent (e.g. saporin, anti-interferon- α , vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten). Bispecific antibodies can be prepared as full length antibodies or antibody fragments [e.g. F(ab')₂ bispecific antibodies].

5 Methods for making bispecific antibodies are known in the art. Traditional production of full length bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein *et al.*, *Nature*, 305:537-539 [1983]). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product
10 yields are low. Similar procedures are disclosed in WO 93/08829, and in Traunecker *et al.*, *EMBO J.*, 10:3655-3659 (1991).

According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2, and
15 CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the
20 construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy
25 chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690. For further details of generating bispecific antibodies see, for example, Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986).

30 According to another approach described in W096/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_H3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size
35 to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

5 Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (US Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in US Patent No. 4,676,980, along with a number of cross-linking techniques.

10 Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, *Science*, 229: 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

15 Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies. Shalaby *et al.*, *J. Exp. Med.*, 175: 217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

20 Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, *J. Immunol.*, 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber *et al.*, *J. Immunol.*, 152:5368 (1994).

35 Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.* *J. Immunol.* 147: 60 (1991).

(vi) *Screening for antibodies with the desired properties*

Techniques for generating antibodies have been described above. Those antibodies having the characteristics described herein are selected.

To select for antibodies which induce cell death, loss of membrane integrity as indicated by, e.g., PI, trypan blue or 7AAD uptake is assessed relative to control. The preferred assay is the "PI uptake assay using BT474 cells". According to this assay, BT474 cells (which can be obtained from the American Type Culture Collection (Rockville, MD)) are cultured in Dulbecco's Modified Eagle Medium (D-MEM):Ham's F-12 (50:50) supplemented with 10% heat-inactivated FBS (Hyclone) and 2 mM L-glutamine. (Thus, the assay is performed in the absence of complement and immune effector cells). The BT474 cells are seeded at a density of 3×10^6 per dish in 100 x 20 mm dishes and allowed to attach overnight. The medium is then removed and replaced with fresh medium alone or medium containing 10 µg/ml of the appropriate MAb. The cells are incubated for a 3 day time period. Following each treatment, monolayers are washed with PBS and detached by trypsinization. Cells are then centrifuged at 1200rpm for 5 minutes at 4°C, the pellet resuspended in 3 ml ice cold Ca^{2+} binding buffer (10 mM Hepes, pH 7.4, 140 mM NaCl, 2.5 mM CaCl_2) and aliquoted into 35 mm strainer-capped 12 x 75 tubes (1ml per tube, 3 tubes per treatment group) for removal of cell clumps. Tubes then receive PI (10 µg/ml). Samples may be analyzed using a FACSCAN™ flow cytometer and FACSCONVERT™ CellQuest software (Becton Dickinson). Those antibodies which induce statistically significant levels of cell death as determined by PI uptake are selected.

In order to select for antibodies which induce apoptosis, an "annexin binding assay using BT474 cells" is available. The BT474 cells are cultured and seeded in dishes as discussed in the preceding paragraph. The medium is then removed and replaced with fresh medium alone or medium containing 10 µg/ml of the MAb. Following a three day incubation period, monolayers are washed with PBS and detached by trypsinization. Cells are then centrifuged, resuspended in Ca^{2+} binding buffer and aliquoted into tubes as discussed above for the cell death assay. Tubes then receive labelled annexin (e.g. annexin V-FTIC) (1 µg/ml). Samples may be analyzed using a FACSCAN™ flow cytometer and FACSCONVERT™ CellQuest software (Becton Dickinson). Those antibodies which induce statistically significant levels of annexin binding relative to control are selected as apoptosis-inducing antibodies.

In addition to the annexin binding assay, a "DNA staining assay using BT474 cells" is available. In order to perform this assay, BT474 cells which have been treated with the antibody of interest as described in the preceding two paragraphs are incubated with 9 µg/ml HOECHST 33342™ for 2 hr at 37°C, then analyzed on an EPICS ELITE™ flow cytometer (Coulter Corporation) using MODFIT LT™ software (Verity Software House). Antibodies which induce a change in the percentage of apoptotic cells which is 2 fold or greater (and preferably 3 fold or greater) than untreated cells (up to 100% apoptotic cells) may be selected as pro-apoptotic antibodies using this assay.

To screen for antibodies which bind to an epitope on ErbB2 bound by an antibody of interest, a routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. Alternatively, epitope mapping can be performed by methods known in the art.

To identify anti-ErbB2 antibodies which inhibit growth of SKBR3 cells in cell culture by 50-100%, the SKBR3 assay described in WO89/06692 can be performed. According to this assay, SKBR3 cells are grown in a 1:1 mixture of F12 and DMEM medium supplemented with 10% fetal bovine serum, glutamine and penicillin-streptomycin. The SKBR3 cells are plated at 20,000 cells in a 35mm cell culture dish (2mls/35mm dish).
 5 2.5µg/ml of the anti-ErbB2 antibody is added per dish. After six days, the number of cells, compared to untreated cells are counted using an electronic COULTER™ cell counter. Those antibodies which inhibit growth of the SKBR3 cells by 50-100% are selected for combination with the apoptotic antibodies as desired.

(vii) *Effector function engineering*

It may be desirable to modify the antibody of the invention with respect to effector function, so as to
 10 enhance the effectiveness of the antibody in treating cancer, for example. For example cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron *et al.*, *J. Exp Med.* 176:1191-1195 (1992) and Shopes, B. *J. Immunol.* 148:2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor
 15 activity may also be prepared using heterobifunctional cross-linkers as described in Wolff *et al. Cancer Research* 53:2560-2565 (1993). Alternatively, an antibody can be engineered which has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al. Anti-Cancer Drug Design* 3:219-230 (1989).

(viii) *Immunoconjugates*

20 The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g. a small molecule toxin or an enzymatically active toxin of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof), or a radioactive isotope (i.e., a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above.
 25 Conjugates of an antibody and one or more small molecule toxins, such as a calicheamicin, a maytansine (U.S. Patent No. 5,208,020), a trichothene, and CC1065 are also contemplated herein. In one preferred embodiment of the invention, the antibody is conjugated to one or more maytansine molecules (e.g. about 1 to about 10 maytansine molecules per antibody molecule). Maytansine may, for example, be converted to May-SS-Me which may be reduced to May-SH3 and reacted with modified antibody (Chari *et al. Cancer Research* 52: 127-
 30 131 [1992]) to generate a maytansinoid-antibody immunoconjugate.

Another immunoconjugate of interest comprises an anti-ErbB2 antibody conjugated to one or more calicheamicin molecules. The calicheamicin family of antibiotics are capable of producing double-stranded DNA breaks at sub-picomolar concentrations. Structural analogues of calicheamicin which may be used include, but are not limited to, γ_1^I , α_2^I , α_3^I , N-acetyl- γ_1^I , PSAG and θ_1^I (Hinman *et al. Cancer Research* 53: 3336-3342 [1993] and
 35 Lode *et al. Cancer Research* 58: 2925-2928 [1998]).

Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolacca americana*

proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcun, croton, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. See, for example, WO 93/21232 published October 28, 1993.

The present invention further contemplates an immunoconjugate formed between an antibody and a compound with nucleolytic activity (e.g. a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase).

A variety of radioactive isotopes are available for the production of radioconjugated anti-ErbB2 antibodies. Examples include At^{211} , I^{131} , I^{125} , Y^{90} , Re^{186} , Re^{188} , Sm^{153} , Bi^{212} , P^{32} and radioactive isotopes of Lu.

Conjugates of the antibody and cytotoxic agent may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate, iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al. Science* 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminopentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026. The linker may be a "cleavable linker" facilitating release of the cytotoxic drug in the cell. For example, an acid-labile linker, peptidase-sensitive linker, dimethyl linker or disulfide-containing linker (Chari *et al. Cancer Research* 52: 127-131 [1992]) may be used.

Alternatively, a fusion protein comprising the anti-ErbB2 antibody and cytotoxic agent may be made, e.g. by recombinant techniques or peptide synthesis.

In yet another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g. avidin) which is conjugated to a cytotoxic agent (e.g. a radionucleotide).

(ix) Immunoliposomes

The anti-ErbB2 antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al., Proc. Natl. Acad. Sci. USA*, 82:3688 (1985); Hwang *et al., Proc. Natl. Acad. Sci. USA*, 77:4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin *et al. J. Biol. Chem.* 257: 286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent or DNA (e.g.

for gene therapy) is optionally contained within the liposome. See Gabizon *et al.* *J. National Cancer Inst.* 81(19)1484 (1989)

(x) *Antibody Dependent Enzyme Mediated Prodrug Therapy (ADEPT)*

The antibodies of the present invention may also be used in ADEPT by conjugating the antibody to a prodrug-activating enzyme which converts a prodrug (*e.g.* a peptidyl chemotherapeutic agent, see WO81/01145) to an active anti-cancer drug. See, for example, WO 88/07378 and U.S. Patent No. 4,975,278.

The enzyme component of the immunoconjugate useful for ADEPT includes any enzyme capable of acting on a prodrug in such a way so as to convert it into its more active, cytotoxic form.

Enzymes that are useful in the method of this invention include, but are not limited to, alkaline phosphatase useful for converting phosphate-containing prodrugs into free drugs; arylsulfatase useful for converting sulfate-containing prodrugs into free drugs; cytosine deaminase useful for converting non-toxic 5-fluorocytosine into the anti-cancer drug, 5-fluorouracil; proteases, such as serratia protease, thermolysin, subtilisin, carboxypeptidases and cathepsins (such as cathepsins B and L), that are useful for converting peptide-containing prodrugs into free drugs; D-alanylcarboxypeptidases, useful for converting prodrugs that contain D-amino acid substituents; carbohydrate-cleaving enzymes such as β -galactosidase and neuraminidase useful for converting glycosylated prodrugs into free drugs; β -lactamase useful for converting drugs derivatized with β -lactams into free drugs; and penicillin amidases, such as penicillin V amidase or penicillin G amidase, useful for converting drugs derivatized at their amine nitrogens with phenoxyacetyl or phenylacetyl groups, respectively, into free drugs. Alternatively, antibodies with enzymatic activity, also known in the art as "abzymes", can be used to convert the prodrugs of the invention into free active drugs (see, *e.g.*, Massey, *Nature* 328: 457-458 [1987]). Antibody-abzyme conjugates can be prepared as described herein for delivery of the abzyme to a tumor cell population.

The enzymes of this invention can be covalently bound to the anti-ErbB2 antibodies by techniques well known in the art such as the use of the heterobifunctional crosslinking reagents discussed above. Alternatively, fusion proteins comprising at least the antigen binding region of an antibody of the invention linked to at least a functionally active portion of an enzyme of the invention can be constructed using recombinant DNA techniques well known in the art (see, *e.g.*, Neuberger *et al.*, *Nature*, 312: 604-608 [1984]).

(xi) *Antibody-salvage receptor binding epitope fusions*

In certain embodiments of the invention, it may be desirable to use an antibody fragment, rather than an intact antibody, to increase tumor penetration, for example. In this case, it may be desirable to modify the antibody fragment in order to increase its serum half life. This may be achieved, for example, by incorporation of a salvage receptor binding epitope into the antibody fragment (*e.g.* by mutation of the appropriate region in the antibody fragment or by incorporating the epitope into a peptide tag that is then fused to the antibody fragment at either end or in the middle, *e.g.*, by DNA or peptide synthesis).

A systematic method for preparing such an antibody variant having an increased *in vivo* half-life comprises several steps. The first involves identifying the sequence and conformation of a salvage receptor binding epitope of an Fc region of an IgG molecule. Once this epitope is identified, the sequence of the antibody of interest is modified to include the sequence and conformation of the identified binding epitope. After the sequence is mutated, the antibody variant is tested to see if it has a longer *in vivo* half-life than that of the original

antibody. If the antibody variant does not have a longer *in vivo* half-life upon testing, its sequence is further altered to include the sequence and conformation of the identified binding epitope. The altered antibody is tested for longer *in vivo* half-life, and this process is continued until a molecule is obtained that exhibits a longer *in vivo* half-life.

5 The salvage receptor binding epitope being thus incorporated into the antibody of interest is any suitable such epitope as defined above, and its nature will depend, *e.g.*, on the type of antibody being modified. The transfer is made such that the antibody of interest still possesses the biological activities described herein.

The epitope preferably constitutes a region wherein any one or more amino acid residues from one or two loops of a Fc domain are transferred to an analogous position of the antibody fragment. Even more preferably, three
10 or more residues from one or two loops of the Fc domain are transferred. Still more preferred, the epitope is taken from the CH2 domain of the Fc region (*e.g.*, of an IgG) and transferred to the CH1, CH3, or V_H region, or more than one such region, of the antibody. Alternatively, the epitope is taken from the CH2 domain of the Fc region and transferred to the C_L region or V_L region, or both, of the antibody fragment. See, US Patent 5,739,277 issued April 14, 1998, expressly incorporated herein by reference.

15 III. Pharmaceutical Formulations

Therapeutic formulations of the antibodies used in accordance with the present invention are prepared for storage by mixing an antibody having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. [1980]), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic
20 to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins,
25 such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.* Zn-protein complexes); and/or non-ionic surfactants such as TWEENTM, PLURONICSTM or polyethylene glycol (PEG).

30 The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to further provide antibodies which bind to EGFR, ErbB2 (*e.g.* an antibody which binds a different epitope on ErbB2), ErbB3, ErbB4, or vascular endothelial factor (VEGF) in the one formulation. Alternatively, or additionally, the composition may comprise a cytotoxic agent, cytokine or growth inhibitory agent.
35 Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes,

albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. [1980].

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

5 Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.* films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, 10 degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss 15 of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

20 The anti-ErbB2 antibody conjugated to a biodegradable nanoparticle (*e.g.* polylactic-co-glycolic acid) to increase tumor-specificity is also contemplated herein.

IV. Treatment with the Anti-ErbB2 Antibodies

The invention herein provides a three-step method for treating a human patient susceptible to or diagnosed with a tumor (or tumors) in which ErbB2 protein is expressed. Generally, the tumor to be treated is a primary tumor. 25 In the first step, a therapeutically effective amount of an anti-ErbB2 antibody is administered to the patient in order to reduce the size of, or eliminate, the tumor (or tumors) in the patient prior to surgery. The patient is optionally further treated with one or more chemotherapeutic agents prior to surgery. In the second step, the tumor is surgically removed according to standard surgical procedures (*e.g.* lumpectomy or mastectomy). Following surgery, in the third step, a therapeutically effective amount of an anti-ErbB2 antibody, or of at least one chemotherapeutic agent, 30 is administered to the patient in order to reduce the likelihood of disease recurrence. Generally, an anti-ErbB2 antibody will be administered to the patient following surgery and, optionally, one or more chemotherapeutic agents will further be administered to the patient during this phase of the therapy.

It is contemplated that, according to the present invention, the anti-ErbB2 antibodies may be used to treat a tumor that expresses, and preferably overexpresses, ErbB2 protein. Exemplary conditions or disorders to be 35 treated herein include benign or malignant tumors (*e.g.* renal, liver, kidney, bladder, breast, gastric, ovarian, colorectal, prostate, pancreatic, lung, vulval, thyroid, hepatic carcinomas; sarcomas; glioblastomas; and various head and neck tumors); leukemias and lymphoid malignancies; other disorders such as neuronal, glial, astrocytal,

hypothalamic and other glandular, macrophagal, epithelial, stromal and blastocoelic disorders; and inflammatory, angiogenic and immunologic disorders.

The antibodies of the invention are administered to a human patient, in accord with known methods, such as intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. Intravenous administration of the antibody is preferred.

Where the anti-ErbB2 antibody is combined with a chemotherapeutic agent, the chemotherapeutic agent is preferably a taxoid, *e.g.* paclitaxel or doxorubicin. Combined administration herein includes coadministration, using separate formulations or a single pharmaceutical formulation, and consecutive administration in either order, wherein preferably there is a time period while both (or all) active agents simultaneously exert their biological activities. Preparation and dosing schedules for such chemotherapeutic agents may be used according to manufacturers' instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such chemotherapy are also described in *Chemotherapy Service Ed.*, M.C. Perry, Williams & Wilkins, Baltimore, MD (1992). Administration of the chemotherapeutic agent may precede, or follow, administration of the antibody or may be given simultaneously therewith. The antibody may be combined with an anti-estrogen compound such as tamoxifen or an anti-progesterone such as onapristone (see, EP 616812) in dosages known for such molecules.

It may be desirable to also administer antibodies against other tumor associated antigens, such as antibodies which bind to the EGFR, ErbB3, ErbB4, or vascular endothelial factor (VEGF). Alternatively, or additionally, two or more anti-ErbB2 antibodies may be co-administered to the patient. Sometimes, it may be beneficial to also administer one or more cytokines to the patient. In a preferred embodiment, the ErbB2 antibody is co-administered with a growth inhibitory agent. For example, the growth inhibitory agent may be administered first, followed by the ErbB2 antibody. However, simultaneous administration or administration of the ErbB2 antibody first is also contemplated. Suitable dosages for the growth inhibitory agent are those presently used and may be lowered due to the combined action (synergy) of the growth inhibitory agent and anti-ErbB2 antibody. Where cardiotoxicity is observed, a cardioprotectant may be administered to the patient as appropriate.

For the prevention or treatment of disease, the appropriate dosage of antibody will depend on the type of disease to be treated, as defined above, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments.

Depending on the type and severity of the disease, about 1 $\mu\text{g/kg}$ to 15 mg/kg (*e.g.* 0.1-20 mg/kg) of antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily dosage might range from about 1 $\mu\text{g/kg}$ to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

V. Articles of Manufacture

In another embodiment of the invention, an article of manufacture containing materials useful for the treatment of the disorders described above is provided. The article of manufacture comprises a container and a label or package insert. Suitable containers include, for example, bottles, vials, syringes, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is effective for treating the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is an anti-ErbB2 antibody. The label or package insert on, or associated with, the container indicates that the composition is used for treating the condition of choice and further indicates treatment of the patient according to the protocol described herein. The article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

Deposit of Materials

The following hybridoma cell lines have been deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, USA (ATCC):

Antibody Designation	ATCC No.	Deposit Date
7C2	ATCC HB-12215	October 17, 1996
7F3	ATCC HB-12216	October 17, 1996
4D5	ATCC CRL 10463	May 24, 1990

Further details of the invention are illustrated by the following non-limiting Example. The disclosures of all citations in the specification are expressly incorporated herein by reference.

Example 1

The anti-ErbB2 IgG₁κ murine monoclonal antibody 4D5, specific for the extracellular domain of ErbB2, was produced as described in Fendly *et al.*, *Cancer Research* 50:1550-1558 (1990) and US Patent 5,677,171 issued October 14, 1997. Briefly, NIH 3T3/HER2-3₄₀₀ cells (expressing approximately 1×10^5 ErbB2 molecules/cell) produced as described in Hudziak *et al. Proc. Natl. Acad. Sci. (USA)* 84:7159 (1987) were harvested with phosphate buffered saline (PBS) containing 25mM EDTA and used to immunize BALB/c mice. The mice were given injections i.p. of 10^7 cells in 0.5ml PBS on weeks, 0, 2, 5 and 7. The mice with antisera that immunoprecipitated ³²P-labeled ErbB2 were given i.p. injections of a wheat germ agglutinin-Sepharose (WGA) purified ErbB2 membrane extract on weeks 9 and 13. This was followed by an i.v. injection of 0.1ml of the ErbB2 preparation and the splenocytes were fused with mouse myeloma line X63-Ag8.653. Hybridoma supernatants were screened for ErbB2-binding by ELISA and radioimmunoprecipitation. MOPC-21 (IgG1), (Cappell, Durham, NC), was used as an isotype-matched control.

A humanized version of the murine 4D5 antibody (HERCEPTIN®) was engineered by inserting the complementarity determining regions of the murine 4D5 antibody into the framework of a consensus human immunoglobulin (IgG₁) (Carter *et al.*, *Proc. Natl. Acad. Sci. USA* 89:4285-4289 [1992]; and US Patent No.

5,821,337 issued October 13, 1998). The resulting humanized anti-ErbB2 monoclonal antibody has high affinity for p185^{HER2} (Dissociation constant [K_d]=0.1 nmol/L), markedly inhibits, *in vitro* and in human xenografts, the growth of breast cancer cells that contain high levels of ErbB2, induces antibody dependent cellular cytotoxicity (ADCC), and has been found clinically active, as a single agent, in patients with ErbB2-overexpressing metastatic breast cancers that had received extensive prior therapy.

HERCEPTIN[®] is produced by a genetically engineered Chinese hamster ovary (CHO) cell line, grown in large scale, that secretes the antibody into the culture medium. The antibody is purified from the CHO culture media using chromatographic and filtration methods. Each lot of antibody used is assayed to verify identity, purity, and potency, as well as to meet Food and Drug Administration requirements for sterility and safety.

Patients with primary breast tumor presentation characterized by overexpression of the ErbB2 (HER2) oncogene [2+ to 3+ as determined by immunohistochemistry or fluorescence *in situ* hybridization (FISH)] are treated herein. Tumor expression of ErbB2 can be determined by immunohistochemical analysis, as previously described (Slamon *et al.*, *Science* 235:177-182 [1987]; Slamon *et al.*, *Science* 244:707-712 [1989]), of a set of thin sections prepared from the patient's paraffin-archived tumor blocks. Tumors are considered to overexpress ErbB2 if at least 25% of tumor cells exhibit characteristic membrane staining for ErbB2.

Patients are first treated with HERCEPTIN[®] for 8-24 weeks, optionally in combination with paclitaxel (TAXOL[®]), in order to reduce the size of, or eliminate, the tumor prior to surgery. On day 0, a 4 mg/kg dose of HERCEPTIN[®] is administered intravenously, over a 90-minute period. Beginning on day 7, patients receive weekly administration of 2 mg/kg antibody (i.v.) over a 90-minute period. Patients may further receive paclitaxel (TAXOL[®]). The initial dose of the HERCEPTIN[®] antibody precedes the first cycle of the chemotherapy regimen by 24 hours. Subsequent doses of the antibody are given immediately before chemotherapy administration, if the initial dose of the antibody is well tolerated. If the first dose of the antibody is not well tolerated, subsequent infusions continue to precede chemotherapy administration by 24 hours. Paclitaxel (TAXOL[®]) is given at a dose of 175 mg/m² over 3 hours by intravenous administration. All patients receiving paclitaxel are premedicated with dexamethasone (or its equivalent) 20 mg x 2, administered orally 12 and 6 hours prior to paclitaxel; diphenhydramine (or its equivalent) 50 mg, iv, administered 30 minutes prior to paclitaxel; and dimetidine (or another H₂ blocker) 300 mg, iv, administered 30 minutes prior to paclitaxel. After the above therapy, classical measures of response may be evaluated immediately prior to surgery; *i.e.*, the sum of the products of the cross-dimensional diameter of any tumor nodules under observation.

Following therapy as described above, the tumor is surgically removed according to standard surgical procedures; lumpectomy or mastectomy. Pathological response may be evaluated at this stage.

After surgery, the patient is treated with HERCEPTIN[®], optionally in combination with paclitaxel (TAXOL[®]), in order to reduce the likelihood of disease recurrence. On day 0, a 4 mg/kg dose of HERCEPTIN[®] is administered intravenously, over a 90-minute period. Beginning on day 7, patients receive weekly administration of 2 mg/kg antibody (i.v.) over a 90-minute period. Therapy with HERCEPTIN[®] is continued for one year. Patients may further receive paclitaxel (TAXOL[®]) for 6-24 weeks. The initial dose of the HERCEPTIN[®] antibody precedes the first cycle of the chemotherapy regimen by 24 hours. Subsequent doses of the antibody are given

immediately before chemotherapy administration, if the initial dose of the antibody is well tolerated. If the first dose of the antibody is not well tolerated, subsequent infusions continue to precede chemotherapy administration by 24 hours. Paclitaxel (TAXOL[®]) is given at a dose of 175 mg/m^2 over 3 hours by intravenous administration. All patients receiving paclitaxel are premedicated as described above.

- 5 Patients treated according to the above therapeutic regimen will display improved overall survival and/or reduced time to tumor progression (TTP).

Claims:

1. A method of treating a human patient susceptible to or diagnosed with a tumor in which ErbB2 protein is expressed comprising the following steps, performed sequentially:
 - (a) treating the patient with a therapeutically effective amount of an anti-ErbB2 antibody;
 - 5 (b) surgically removing the tumor; and then
 - (c) treating the patient with a therapeutically effective amount of an anti-ErbB2 antibody or of a chemotherapeutic agent.
- 10 2. The method of claim 1 wherein step (a) further comprises treating the patient with a therapeutically effective amount of a chemotherapeutic agent.
3. The method of claim 1 wherein step (c) comprises treating the patient with a therapeutically effective amount of an anti-ErbB2 antibody.
- 15 4. The method of claim 3 wherein step (c) further comprises treating the patient with a therapeutically effective amount of a chemotherapeutic agent.
5. The method of claim 1 wherein the tumor overexpresses ErbB2 protein.
- 20 6. The method of claim 5 wherein the tumor is selected from the group consisting of a breast tumor, squamous cell tumor, small-cell lung tumor, non-small cell lung tumor, gastrointestinal tumor, pancreatic tumor, glioblastoma, cervical tumor, ovarian tumor, liver tumor, bladder tumor, hepatoma, colon tumor, colorectal tumor, endometrial tumor, salivary gland tumor, kidney tumor, prostate tumor, vulval tumor, thyroid tumor, hepatic tumor, head tumor and neck tumor.
- 25 7. The method of claim 6 wherein the tumor is a breast tumor.
8. The method of claim 1 wherein the antibody binds to the extracellular domain of the ErbB2 protein.
- 30 9. The method of claim 8 wherein the antibody binds to epitope 4D5 within the ErbB2 extracellular domain sequence.
10. The method of claim 9 wherein the antibody is a humanized 4D5 anti-ErbB2 antibody.
- 35 11. The method of claim 2 wherein the chemotherapeutic agent is a taxoid.
12. The method of claim 11 wherein the taxoid is paclitaxel or doxorubicin.

13. The method of claim 4 wherein the chemotherapeutic agent is a taxoid.
 14. An article of manufacture comprising a container, a composition within the container comprising an anti-ErbB2 antibody and a package insert instructing the user of the composition to treat a patient essentially according to the method of claim 1.
- 5


1 / 1

3H4 aa 541-599

4D5 aa 529-625

7C2 aa 22-53

7F3 aa 22-53



3H4 epitope (SEQ ID NO:3)

VEECRVLQGLPREYVNARHCLPCHPECQPQNGSVTCFGPEADQCVACAHYKDPPFCVAR
 | 541 | 599

4D5 epitope (SEQ ID NO:4)

LPCHPECQPQNGSVTCFGPEADQCVACAHYKDPPFCVARCPGKPDLSYMPIWKFPDEEGACQP
 | 561 | 625

FIG. 1

1 MELAALCRWGLLLALLPPGAASTQVCTGTDMLRLPA
 38 SPETHLDMLRHLVYQGCQVVOGNLELTYPNASLSFL
 75 QDIQEVQGYVLIHNVQVRQVPLQRLRIVRGTLQFEDN
 112 YALAVLDNGDPLNNTTPVTGASPGGLRELQLRSLTEI
 149 LKGGVLIQRNPQLCYQDTILWKDIFHKNNQLALTLD
 186 TNRSRA

FIG. 2

SUBSTITUTE SHEET (RULE 26)

Sequence Listing

<110> Genentech, Inc.
 5 <120> TREATMENT WITH ANTI-ErbB2 ANTIBODIES
 <130> P1757R1PCT
 <141> 2000-05-09
 10 <150> US 60/134,085
 <151> 1999-05-14
 <160> 4
 15 <210> 1
 <211> 166
 <212> PRT
 <213> Homo sapiens
 20 <400> 1
 Cys Thr Gly Thr Asp Met Lys Leu Arg Leu Pro Ala Ser Pro Glu
 1 5 10 15
 25 Thr His Leu Asp Met Leu Arg His Leu Tyr Gln Gly Cys Gln Val
 20 25 30
 Val Gln Gly Asn Leu Glu Leu Thr Tyr Leu Pro Thr Asn Ala Ser
 35 40 45
 30 Leu Ser Phe Leu Gln Asp Ile Gln Glu Val Gln Gly Tyr Val Leu
 50 55 60
 Ile Ala His Asn Gln Val Arg Gln Val Pro Leu Gln Arg Leu Arg
 35 65 70 75
 Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr Ala Leu Ala
 80 85 90
 40 Val Leu Asp Asn Gly Asp Pro Leu Asn Asn Thr Thr Pro Val Thr
 95 100 105
 Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser Leu
 110 115 120
 45 Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln
 125 130 135
 Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys
 50 140 145 150
 Asn Asn Gln Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg
 155 160 165
 55 Ala
 <210> 2
 <211> 32
 60 <212> PRT
 <213> Homo sapiens
 <400> 2
 Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys Leu Arg Leu Pro

1 5 10 15
 Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His Leu Tyr Gln
 20 25 30
 5 Gly Cys

 <210> 3
 10 <211> 59
 <212> PRT
 <213> Homo sapiens

 <400> 3
 15 Val Glu Glu Cys Arg Val Leu Gln Gly Leu Pro Arg Glu Tyr Val
 1 5 10 15
 Asn Ala Arg His Cys Leu Pro Cys His Pro Glu Cys Gln Pro Gln
 20 25 30
 20 Asn Gly Ser Val Thr Cys Phe Gly Pro Glu Ala Asp Gln Cys Val
 35 40 45
 Ala Cys Ala His Tyr Lys Asp Pro Pro Phe Cys Val Ala Arg
 25 50 55

 <210> 4
 <211> 65
 <212> PRT
 30 <213> Homo sapiens

 <400> 4
 Leu Pro Cys His Pro Glu Cys Gln Pro Gln Asn Gly Ser Val Thr
 1 5 10 15
 35 Cys Phe Gly Pro Glu Ala Asp Gln Cys Val Ala Cys Ala His Tyr
 20 25 30
 Lys Asp Pro Pro Phe Cys Val Ala Arg Cys Pro Ser Gly Val Lys
 40 35 40 45
 Pro Asp Leu Ser Tyr Met Pro Ile Trp Lys Phe Pro Asp Glu Glu
 50 55 60
 45 Gly Ala Cys Gln Pro
 65

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 00/12552

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K39/395 A61P35/00 //(A61K39/395,31:335)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GOLDENBERG M M: "Trastuzumab, a recombinant DNA-derived humanized monoclonal antibody, a novel agent for the treatment of metastatic breast cancer." CLINICAL THERAPEUTICS, (1999 FEB) 21 (2) 309-18. REF: 28 , XP000918210 page 309 -page 310, paragraph 1 page 314, last paragraph -page 315, right-hand column, paragraph 1 ---	1-14
X	DEES E C ET AL: "Recent advances in systemic therapy for breast cancer." CURRENT OPINION IN ONCOLOGY, (1998 NOV) 10 (6) 517-22. REF: 39 , XP000918214 page 520, right-hand column, paragraph 2 - paragraph 5 --- -/--	1-14



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

18 September 2000

Date of mailing of the international search report

10 OCTOBER 2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl
Fax: (+31-70) 340-3016

Authorized officer

von Ballmoos, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/12552

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BASELGA J ET AL: "Anti HER2 humanized monoclonal antibody (Mab) alone and in combination with chemotherapy against human breast carcinoma xenografts" BREAST CANCER, PROCEEDINGS OF ASCO, VOL. 13 MARCH 1994, ABSTRACT *53, XP000918303 cited in the application the whole document	1-14
A	--- CARTER P ET AL: "HUMANIZATION OF AN ANTI-P185HER2 ANTIBODY FOR HUMAN CANCER THERAPY" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA,US,NATIONAL ACADEMY OF SCIENCE. WASHINGTON, vol. 89, no. 10, 15 May 1992 (1992-05-15), pages 4285-4289, XP000275844 ISSN: 0027-8424 cited in the application the whole document	1-14
P,X	--- WO 99 31140 A (GENENTECH INC) 24 June 1999 (1999-06-24) the whole document -----	1-14

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 00/12552

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1-13 are directed to a method of treatment of the human body, the search has been carried out and based on the alleged effects of the composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

Information on patent family members

PCT/US 00/12552

Form PCT/ISA/210 (patent family annex) (July 1992)